

## Removing the Kid Gloves in Neurologic Rehabilitation

### Reference List

1. Hornby TG, Campbell DD, Kahn JH, Demott T, Moore JL, Roth HR. Enhanced gait-related improvements following therapist- vs. robotic-assisted locomotor training in subjects with chronic stroke: a randomized controlled study. *Stroke*. 2008. 39(6):1786-92. [Link to article](#)
2. Hidler J, Nichols D, Pelliccio M, Brady K, Campbell DD, Kahn JH, Hornby TG. Multicenter randomized clinical trial evaluating the effectiveness of the Lokomat in subacute stroke. *Neurorehabil Neural Repair*. 2009 Jan;23(1):5-13. [Link to article](#)
3. Moore JL, Roth EJ, Killian C, Hornby TG. Locomotor training improves daily stepping activity and gait efficiency in individuals poststroke who have reached a "plateau" in recovery. *Stroke*. 2010 Jan;41(1):129-35 [Link to article](#)
4. Duncan PW, Sullivan KJ, Behrman AL, Azen SP, Wu SS, Nadeau SE, Dobkin BH, Rose DK, Tilson JK, Cen S, Hayden SK; LEAPS Investigative Team. Body-weight-supported treadmill rehabilitation after stroke. *N Engl J Med*. 2011 May 26;364(21):2026-36. doi: 10.1056/NEJMoa1010790. [Link to article](#)
5. Holleran CL, Rodriguez KS, Echaz A, Leech KA, Hornby TG. Potential contributions of training intensity on locomotor performance in individuals with chronic stroke. *J Neurol Phys Ther*. 2015 Apr;39(2):95-102. [Download article](#)
6. Brazg G, Fahey M., Holleran CL., Connolly M, Woodward J., Hennessy PW., Schmit, BD., Hornby TG. Effects of training intensity on locomotor performance in individuals with chronic spinal cord injury: a randomized crossover study. *NRR* 2017 Oct-Nov;31(10-11):944-954. [Link to article](#)
7. Holleran CL, Straube DD, Kinnaird CR, Leddy AL, Hornby TG. Feasibility and potential efficacy of high-intensity stepping training in variable contexts in subacute and chronic stroke. *Neurorehabil Neural Repair*. 2014 Sep;28(7):643-51. *Note: Protocol is included in this article appendix* [Download article](#)
8. Straube DD, Holleran CL, Kinnaird CR, Leddy AL, Hennessy PW, Hornby TG. Effects of dynamic stepping training on nonlocomotor tasks in individuals poststroke. *Phys Ther*. 2014 Jul;94(7):921-33. [Link to article](#)
9. Hornby TG, Holleran CL, Hennessy PW, Leddy AL, Connolly M, Camardo J, Woodward J, Mahtani G, Lovell L, Roth EJ. Variable Intensive Early Walking Poststroke (VIEWS): A Randomized Controlled Trial. *Neurorehabil Neural Repair*. 2015 Sep 3. [Link to article](#)
10. Hornby TG, Henderson CE, Plawecki A, Lucas E, Lotter J, Holthus M, Brazg G, Fahey M, Woodward J, Ardestani M, Roth EJ. Contributions of Stepping Intensity and Variability to Mobility in Individuals Poststroke. *Stroke*. 2019 Sep;50(9):2492-2499. [Link to article](#)
11. Leddy AL, Connolly M, Holleran CL, Hennessy PW, Woodward J, Arena RA, Roth EJ, Hornby TG. Alterations in Aerobic Exercise Performance and Gait Economy Following High-Intensity Dynamic Stepping Training in Persons With Subacute Stroke. *J Neurol Phys Ther*. 2016 Oct;40(4):239-48. [Link to article](#)

12. Pang MY, Charlesworth SA, Lau RW, Chung RC. Using aerobic exercise to improve health outcomes and quality of life in stroke: evidence-based exercise prescription recommendations. *Cerebrovasc Dis*. 2013;35:7-22. [Link to article](#)
13. Mahtani GB, Kinnaird CR, Connolly M, Holleran CL, Hennessy PW, Woodward J, Brazg G, Roth EJ, Hornby TG. Altered Sagittal- and Frontal-Plane Kinematics Following High-Intensity Stepping Training Versus Conventional Interventions in Subacute Stroke. *Phys Ther*. 2016 Sep 15. [Link to article](#)
14. Tyrell CM, Roos MA, Rudolph KS, Reisman DS. Influence of systematic increases in treadmill walking speed on gait kinematics after stroke. *Phys Ther*. 2011 Mar;91(3):392-403. [Link to article](#)
15. Reisman DS, Wityk R, Silver K, Bastian AJ. Locomotor adaptation on a split-belt treadmill can improve walking symmetry post-stroke. *Brain*. 2007 Jul;130(Pt 7):1861-72. [Link to article](#)
16. Ardestani MM, Kinnaird CR, Henderson CE, Hornby TG. Compensation or Recovery? Altered Kinetics and Neuromuscular Synergies Following High-Intensity Stepping Training Poststroke. *Neurorehabil Neural Repair*. 2019 Jan;33(1):47-58. [Link to article](#)
17. Ardestani MM, Henderson CE, Salehi SH, Mahtani GB, Schmit BD, Hornby TG. Kinematic and Neuromuscular Adaptations in Incomplete Spinal Cord Injury after High- versus Low-Intensity Locomotor Training. *J Neurotrauma*. 2019 Feb 1. [Link to article](#)
18. Ardestani MM, Henderson CE, Mahtani G, Connolly M, Hornby TG. Locomotor Kinematics and Kinetics Following High-Intensity Stepping Training in Variable Contexts Poststroke. *Neurorehabil Neural Repair*. 2020 Jul;34(7):652-660. doi:
19. Hornby TG, Holleran CL, Leddy AL, et al. Feasibility of Focused Stepping Practice During Inpatient Rehabilitation Poststroke and Potential Contributions to Mobility Outcomes. *Neurorehabil Neural Repair*. 2015;29(10):923-32. [Link to article](#)
20. Amy J Bastian. Learning to predict the future: the cerebellum adapts feedforward movement control. *Current Opinion in Neurobiology* 2006, 16:645–649. [Link to abstract](#)
21. Darcy S. Reisman, PhD, Heather McLean, Jennifer Keller, Kelly A. Danks, and Amy J. Bastian. Repeated Split-Belt Treadmill Training Improves Poststroke Step Length Asymmetry. *Neurorehabilitation and Neural Repair* 27(5) 460–468. [Link to article](#)
22. Moore JL, Nordvik JE, Erichsen A, Rosseland I, Bø E, Hornby TG; FIRST-Oslo Team. Implementation of High-Intensity Stepping Training During Inpatient Stroke Rehabilitation Improves Functional Outcomes. *Stroke*. 2019 Dec 30:STROKEAHA119027450. [Link to abstract](#)
23. Hornby TG, Reisman DS, Ward IG, Scheets PL, Miller A, Haddad D, Fox EJ, Fritz NE, Hawkins K, Henderson CE, Hendron KL, Holleran CL, Lynskey JE, Walter A; and the Locomotor CPG Appraisal Team. Clinical Practice Guideline to Improve Locomotor Function Following Chronic Stroke, Incomplete Spinal Cord Injury, and Brain Injury. *J Neurol Phys Ther*. 2020 Jan;44(1):49-100. [Link to article](#)

24. Hornby TG, Straube DS, Kinnaird CR, Holleran CL, Echaz AJ, Rodriguez KS, Wagner EJ, Narducci EA. Importance of Specificity, Amount, and Intensity of Locomotor Training to Improve Ambulatory Function in Patients Poststroke. *Top Stroke Rehabil.* 2011. Jul-Aug;18(4):293-307. [Download article](#)
25. Leech KA, Hornby TG. High-Intensity Locomotor Exercise Increases Brain-Derived Neurotrophic Factor in Individuals with Incomplete Spinal Cord Injury. *J* [Download article](#)
26. Woodward JL, Connolly M, Hennessy PW, Holleran CL, Mahtani GB, Brazg G, Fahey M, Maganti K, Hornby TG. Cardiopulmonary Responses During Clinical Laboratory Gait Assessments in People with Chronic Stroke. *Phys Ther.* 2019 Jan 1;99(1):86-97. [Download article](#)
27. Lewek MD, Cruz TH, Moore JL, Roth HR, Dhaher YY, Hornby TG. Allowing intralimb kinematic variability Allowing intralimb kinematic variability during locomotor training poststroke improves kinematic consistency: a subgroup analysis from a randomized clinical trial. *Phys Ther.* 2009;89(8):829-839. [Download article](#)
28. Lotter JK, Henderson CE, Plawecki A, Holthus ME, Lucas EH, Ardestani MM, Schmit BD, Hornby TG. Task-Specific Versus Impairment-Based Training on Locomotor Performance in Individuals With Chronic Spinal Cord Injury: A Randomized Crossover Study. *Neurorehabil Neural Repair.* 2020 Jul;34(7):627-639. [Download article](#)
29. Hornby TG, Henderson CE, Plawecki A, Lucas E, Lotter J, Holthus M, Brazg G, Fahey M, Woodward J, Ardestani M, Roth EJ. Contributions of Stepping Intensity and Variability to Mobility in Individuals Poststroke. *Stroke.* 2019 Sep;50(9):2492-2499. [Download article](#)